



# FERTILIZER W/ Barricade® 0.3%

887238

0-0-7

FOR PRE-EMERGENCE CONTROL OF WEED GRASSES AND BROADLEAF WEEDS IN ESTABLISHED TURF GRASSES, LAWNS, GOLF COURSE FAIRWAYS AND SOD FARMS, (NOT FOR PUTTING GREENS). USE FOR PRE-EMERGENCE WEED CONTROL AROUND LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWER PLANTINGS.

**Active Ingredient:**

Prodiamine\*\*\*..... 0.300%

**Other Ingredients:**..... 99.700%

Total..... 100.000%

\*\*\*CAS 29091-21-2

EPA Reg. No. 52287-20

EPA Est. No.52287-AL-001

**GUARANTEED ANALYSIS**

**Soluble Potash (K<sub>2</sub>O)**..... **7.0000%**

**Derived From: Muriate of Potash**

**Chlorine (Cl), Not more than**..... **5.4190%**

F352 Density - 88 lb./cu. ft.

Batch No. 2108-0614

**Net Wt. 50 LBS. (22.7 KG)**

## KEEP OUT OF REACH OF CHILDREN

### CAUTION

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

**READ ENTIRE LABEL BEFORE USING**

**FIRST AID**

If swallowed:	<ul style="list-style-type: none"> <li>Call a poison control center or doctor immediately for treatment advice</li> <li>Have person sip a glass of water if able to swallow</li> <li>Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>Do not give anything by mouth to an unconscious person.</li> </ul>
If in eyes:	<ul style="list-style-type: none"> <li>Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If on skin:	<ul style="list-style-type: none"> <li>Take off contaminated clothing.</li> <li>Rinse skin immediately with plenty of water for 15-20 minutes.</li> <li>Call a poison control center or doctor for treatment advice.</li> </ul>
If inhaled:	<ul style="list-style-type: none"> <li>Move person to fresh air.</li> <li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible.</li> <li>Call a poison control center or doctor for further treatment advice.</li> </ul>

Have the product container or label with you when calling a poison control center, or doctor, or when going for treatment.

**PRECAUTIONARY STATEMENTS**

**HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequent repeated skin contact while handling the material may cause allergic reactions in some individuals.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

Applicators and other handlers must wear long sleeved shirt and long pants, chemical resistant gloves made of butyl rubber >14 mils or natural rubber > 14 mils or natural rubber > 14 mils or nitrile rubber > 14 mils, shoes plus socks and a dust mask. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**User Safety Recommendations**

Users should: Wash hands thoroughly after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product has low solubility in water. At limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated area may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. This product is a turf fertilizer and a preemergence herbicide that provides residual control of many grass and broadleaf weeds in established turf grasses, golf courses (excluding putting greens), for sod farms, and lawns, and around landscape ornamentals, established perennials and wildflower plantings. It controls susceptible weeds by inhibiting weed seed germination and root development. **Do not graze or feed livestock forage cut from areas treated with this product. Do not apply aerially. Do not apply to golf course putting greens. Failure to follow the directions for use and precautions on this label may result in poor weed control, crop injury, or illegal residues.**

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard (sod farms, commercial nurseries). Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is: coveralls; chemical resistant gloves made of butyl rubber > 14 mils or natural rubber > 14mils or nitrile rubber > 14 mils; shoes plus socks.

**NON-AGRUCULTURAL USE REQUIREMENTS**

The requirements of this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR part 170). The "WPSP" applies when this product is used to produce agricultural plants on farms, forests or nurseries. Applicators and handlers should wear long-sleeved shirt, long pants, chemical resistant gloves, shoes with socks and protective eyewear. Thoroughly wash face and hands with soap and water after exposure and before eating or smoking. Launder exposed clothing before reuse. Do not allow reentry to treated areas until dusts have settled and the turfgrass or soil is dry.

**APPLICATION DIRECTIONS**

Apply uniformly with suitable, calibrated application equipment. The size and bulk density of granules in this product may vary; therefore, calibrate your granular applicator prior to application. A calibrated applicator must be used. Follow directions in the owner's manual for calibrating your applicator.

**RATES OF APPLICATION**

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. All applications must be made prior to germination of the target weeds. This product will not control established weeds. The length of time of residual pre-emergence weed control provided by this product is related to the rate applied. The maximum use rate selection must be based on turf species.

## WEEDS CONTROLLED

When used in accordance with this label this product will provide control of the following weeds:		
Barnyardgrass	Henbit	Purslane, Common
Bluegrass, Annual ( <i>Poa Annua</i> )	Itchgrass	Pusley, Florida
Carpetweed	Johnsongrass (from seed)	Rescuegrass <sup>(3)</sup>
Chickweed, Common	Junglerice	Shepherd's Purse <sup>(2)</sup>
Chickweed, Mouseear (from seed)	Knotweed	Signalgrass, Broadleaf
Crabgrass (Large, smooth)	Kochia	Speedwell, Persian
Crowfootgrass	Lambsquarter, Common	Sprangletop
Cupgrass, Woolly	Lovegrass	Spurge, Prostrate
Foxtails, Annual	Panicum (Texas, Fall, Browntop)	Witchgrass
Goosegrass <sup>(1)</sup>	Pigweed	Woodsorrel, Yellow (from seed)

(1) In many areas a single application of 215 to 500 lbs. per acre will control Goosegrass. However, under heavy Goosegrass pressure and/or an extended growing season, most effective weed control may be obtained by making an initial application of 215 to 334 lbs per acre followed after 60 to 90 days by a second application at doses that would not exceed those given in the [Maximum Annual Rates](#) section. Do not exceed the maximum rates in the [Maximum Annual Rates](#) section. (2) Application for Shepherd's Purse should be made in late summer, fall or winter prior to germination. (3) Suppression only of Rescuegrass.

Sequential application may be made so long as the total amount of product applied does not exceed the maximum annual application rates. All applications must be made prior to germination of the weed seeds.

## MAXIMUM ANNUAL RATES

This product is for use on the turf grass species listed below and on landscape ornamental species as listed in the Tolerant Ornamental Species section of this label. Do not exceed the maximum yearly rate as given in the following table:

TURF GRASS		
MAXIMUM APPLICATION RATE/CALENDAR YEAR OF PRODUCT BY TURF GRASS SPECIES <sup>(1)</sup>		
Turf Species	Lbs. product/A	Lbs. a.i./A
Creeping Bentgrass	215 lbs.	0.65 lbs.
Creeping Red Fescue	250 lbs.	0.75 lbs.
Buffalograss	334 lbs.	1.00 lbs.
Kentucky Bluegrass	334 lbs.	1.00 lbs.
Perennial Ryegrass	334 lbs.	1.00 lbs.
Bermudagrass <sup>(2)</sup>	500 lbs.	1.50 lbs.
Bahiagrass	500 lbs.	1.50 lbs.
Centipedegrass	500 lbs.	1.50 lbs.
Seashore Paspalum	500 lbs.	1.50 lbs.
St. Augustinegrass	500 lbs.	1.50 lbs.
Tall Fescue (including turf type)	500 lbs.	1.50 lbs.
Zoysia	500 lbs.	1.50 lbs.
LANDSCAPE ORNAMENTALS		
Tolerant Ornamental Species	500 lbs.	1.50 lbs.

(1) These are the maximum rates per calendar year by species limitations. (2) May be used on newly-sprigged or plugged bermudagrass at rates not to exceed 173 lbs. per acre. Newly-sprigged or plugged bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced product rates used in sprigging situations. Do not apply more than 500 lbs. of this product to the same area per calendar year. Use higher rates of product to achieve higher levels of fertility and longer periods of weed control for each turf type, but do not exceed the maximum application rates specified in the [Maximum Annual Rates](#) section.

## ESTABLISHED TURF

This product is a pre-emergence herbicide that, when properly applied, will control certain grass and broadleaf weeds listed in the Weeds Controlled Table in established turf grass and lawns. The maximum amount of material that may be applied per year is given for each turf grass species in the [Maximum Annual Rates](#) section of this label. Most effective weed control

in turf grasses will be obtained when this product is activated by at least 1/4 inch of rainfall or irrigation prior to weed seed germination and within 14 days following application. See the map for approximate crabgrass seed germination dates.



- 1: Mid-north & northwestern.....After May 30
- 2: Upper/mid-states & northeast.....After May 10
- 3: Lower mid-states.....After April 20
- 4: Upper southern states.....After March 20
- 5: Warm season states.....January 1 to March 20

## Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns:

(1) Application of this product may thin emerged annual bluegrass and newly overseeded grasses. (2) Do not apply to overseeded turf within 60 days after seeding or until after the second mowing whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil. (3) Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. (4) Disturbing the herbicide barrier with cultural practices such as disking may result in reduced weed control.

## Use Restrictions

(1) Do not cut (harvest) treated sod before 120 days after application. Do not apply to newly set sod until the following year. (2) Do not apply this product to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species.

## WHEN TO APPLY AFTER OVERSEEDING TURF

Do not apply to overseeded turf within 60 days after seeding or until after the second mowing, whichever is longer. Injury to desirable seedlings is likely if this product is applied before seedling secondary roots are in the second inch of soil, not thatch plus soil.

## WHEN TO OVERSEED AFTER APPLICATION

This product will inhibit the germination of turf species if overseeded too soon after application. Follow rates and intervals in the table below for best overseeding and/or reseeding results.

### Months Before Overseeding\*

Product Rate Lbs./A	Lbs. ai/Acre	North	Transition*	South
167	0.50	4	4	4
215	0.65	5	4	4
250	0.75	6	5	5
267	0.80	-	6	6
334	1.00	-	7	7
380	1.14	-	-	9
433	1.30	-	-	10
500	1.50	-	-	12

\*Transition states are: DE, MD, VA, WV, KY, MO, KS; North states are those north of transition states; South states are those south of transition states.

## LANDSCAPE ORNAMENTALS INCLUDING ESTABLISHED PERENNIALS AND WILD FLOWER PLANTINGS

This product may be applied for residual pre-emergence weed control in ornamentals listed on this label.

## USE RATES

Apply at 215 to 500 lbs. per acre of landscape plants, in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 500 lbs. per acre.

## APPLICATION TIMING AND INFORMATION

Apply uniformly to the surface of the soil in ornamental planting beds with a granular applicator.

This product is a pre-emergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 1/4 inch of rainfall or irrigation, or with shallow (1 to 2 inches) mechanical incorporation, prior to weed seed germination and within 14 days following application. Best weed control is obtained when this product is applied to soil free of clods, weeds, and debris such as leaves. Prior to application of this product, control existing vegetation by hand weeding, cultivation, or the use of an appropriate post-emergence herbicide.

## ESTABLISHED LANDSCAPE ORNAMENTALS

This product may be applied to established ornamentals as a broadcast, over the top, or directed application. Irrigation or rainfall soon after application will wash residues off plant foliage and activate this product in the soil.

## NEWLY TRANSPLANTED ORNAMENTALS

This product may be applied to newly transplanted ornamentals as a broadcast, over the top, or directed application. Delay application to allow soil to settle around new transplants and water thoroughly before applying this product. Apply after cuttings from roots have become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union. Irrigate after application as above directed.

## TOLERANT ORNAMENTAL SPECIES

This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when this product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought, or frost damage) some cultivars of listed plants may be sensitive to this product.

## FOR USE AROUND LANDSCAPE ORNAMENTALS, INCLUDING ESTABLISHED PERENNIALS AND WILDFLOWERS, AS LISTED.

Scientific Name	Common name
Abies spp.	Fir Species**
Abelia grandiflora	Abelia: Sherwood
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Achillea, sp.	Yarrow: King Edward
Actinidia chinensis*	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Agapanthus orientalis	
Akebia quintata	Five-Leaf or Chocolate Vine
Allium cernuum	Lady's Leek, Nodding Onion
Anemone hybrid	Japanese Anemone
Aquilegia sp.	Aquilegia: Red and Gold
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Artemisia sp.	Wormwood: Silver Mound, Castle
Aster sp.	Aster: Bonny Blue, Purple Dome
Aster X Frikartii	
Athrium Filiz-femina	Lady Fern: Fern Lady
Aucuba japonica	Japanese Aucuba
Begonia sp.	Fibrous Begonia: Hardy Grandis
Berberis gladywnensis	Barberry
B. julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
B. verruculosa	Warty Barberry
Bergenia cordifolia	
Boltonia asteroides	Snowbank
Bougainvillea spp.	Bougainvillea
Buddleia davidii	Butterfly-Bush (Dwarf Blue); Royal Red
Buxus microphylla	Japanese Boxwood

Scientific Name	Common name
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Campanula carpatica	Tussock Bellflower (White Clips)
Campis X Tagliabuana	Trumpet Creeper, Trumpet Flower, Madame Galen
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Ceratostigma Plumbagonoides	
Chamaecyparis pisifera	False Cypress
Chrysanthemum nipponicum	
Cleyera japonica	Cleyera
Citrus spp.*	Citrus species*
Coreopsis sp.	Coreopsis (Calliopsis): Early Sunrise, Moonbeam
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotonneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
Cretaegus spp.	Hawthorne
Crocsmia sp.	Lucifer
Cupressus sempervirens	Italian Cypress
Delosperma alba	White Trailing Ice Plant
Delosperma sp.	Cooperi Pink
Delphinium sp.	Larkspur: Blue Elf
Dianthus deltoides	Dianthus: Maiden Pinks "Zing"
D. gratianopolitanus	Cheddar Pink
Dodonea viscosa	Hop Bush
Echinacea purpurea	Coneflower, Purple; Mangus
Elaeagnus pungens	Silverberry
Euonymus fortune	Wintercreeper
E. japonica	Japanese Spindle Tree (Evergreen Euonymus)
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
F. suspense	Weeping Forsythia
F. viridissima	Greenstem Forsythia
Gaillardia sp.	Gaillardia, Blanket Flower: "Goblin"
Gardenia jasminoides	Gardenia, Cape-Jasmine, Radicans
Gaura sp.	
Gentiana dahurica	Gentian
Geranium cinereum	Cranesbill
Gladiolus spp.**	Gladiolus species**
Gypsophila repens	Baby's Breath
Hedera helix	English Ivy
Helianthemum sp.	Sunrose
Hemerocallis sp.	Daylily: Aztec Gold, Stella De Oro, Tender Love
Hibiscus**	Rose of Sharon**
Hibiscus sp.	Mallow: Disco Belle White
Hibiscus rosa-sinensis	Chinese Hibiscus
Hosta plantaginea	Hosta, Plantain Lily (Fragrant)
H. sieboldiana	Hosta, "Searsucker"
Houttuynia cordata var. Variegata	
Hydrangea macrophylla	Bigleaf hydrangea
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly; Helliery
I. opaca	American Holly
I. pernyi	Holly
I. vomitoria	Yaupon Holly; Schillings
Inula ensifolia	
Iris ensata	Sword-Leaved Iris; Jodlesong
Iris siberica	Siberian Iris; Cabernet
Iris spp.**	Iris species**
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper; Nick's Compact, Parsonii
J. conferta	Shore Juniper; Blue Pacific

Scientific Name	Common name
J. davurica	Parsoni
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crape Myrtle
Lagerstromia indica X Fauriei	Crape Myrtle; Tuscarora
Lantana montevidensis	Weeping Lantana
Lavender sp.	Lavender; Munstead
Leontopodium alpinum	Edelweiss
Ligustrum sinense	Chinese Privet; Variegata
L. amurensis	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Lillium sp.	Lily; Jazz
Liriope muscari	Big Blue Lily; Turf; Silver Mound; Evergreen Giant
Liriope Muscari var. Variegata	Liriope, Variegated
Lirope spicata	Liriope, Creeping
Lobelia Cardinalis	Cardinal Flower, Indian Pink
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Loropetalum Chinese	Burgundy
Lythrum sp.	Loosestrife; Modern Pink
Magnolia spp.**	Magnolia species**
Maleophora luteola	Ice Plant
Malus sp.*	Crabapple*
Miscanthus sinensis	Yaku Jima, Silberfeder
Nandina domestica	Heavenly Bamboo
Narcissus spp.**	Narcissus species**
Nerium oleander	Oleander
Olea europea*	Olive*
Ophiopogon japonicus	Mondo Grass
Osmanthus heterophyllus	Osmanthus (False Holly); Gulf tide
Osteospermum fruticosum	Trailing African Daisy
Oxydendrum arboretum	Sourwood
Paeonia suffruticosa	Tree Peony
Pennisetum setaceum	Fountain Grass (Dwarf)
Perovskia atriplicifolia	
Persea Americana	Avocado*
Photinia fraseri	Frasier's Photinia ((Redtip)
Physotegia Virginiana	Dragonhead, False; Vivid
Picca spp.**	Spruce species** (Colorado Blue, Norway, etc.)
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary island Pine
P.elliottii	Slash Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. palustris	Longleaf Pine
P. radiata	Monterey Pine
P. thunbergiana	Japanese Black Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
P. taeda	Loblolly Pine
P. virginiana	Virginia Pine
Pistachio sp.*	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum
P. tobira	Japanese Pittosporum
Posocarpus Macrophyllus	Japanese Yew
Prunus fauocerasus	English Laurel
Prunus sp.	Almond, Apricot, Nectarine, Peach, Plum and Prune*
Pseudotsuga menziesii**	Douglas Fir**
Pyracantha Coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus Sp.	Bradford Pear sp.
Quercus Shumardii	Oak, Shumard's Red
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne

Scientific Name	Common name
Raphiolepis umbellata	Yedda Hawthorn
Rhododendron (including Azalea)	'Coro Bells' 'Delaware, Valley White', 'Flame Creeper', 'formosa', 'Girard Crimson', George L. Tabor', 'Hino-crimson', 'PJM', 'Roseum Elegans', 'Wakeiebisu', 'White Gumbo'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis*	Rosemary*
Rudbeckia sp.	Black-Eyed susan; Goldstrum
Santolina, virens	Santolina
Saxifraga sp.	Axifrage; Purple Dome
Scabiosa sp.	Pincushion Flower
Sedum album	Stonecrop
Sedum cauticola	Stonecrop; Lidakense
Sedum dasyphyllum	Stonecrop
Sedum spurium	Sonecrop; Dragon's Blood
Spiraea Bumalda	Spirea; Anthony Waterer
Syzgium paniculatum	Australian Brushcherry
Taxus cuspidate	Japanese Yew
T. media	Yew
Teucrium sp.	Germander
Thalictrum dipterocarpum	Meadow Rue
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga Canadensis	Canada Hemlock
Tulipa spp.	Tulip species
Veronica sp.	Veronica, Speedwell; Sunny Border
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rigidum	Canary Island Viburnum
V. suspensum	Arrowood Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca minor	Common Periwinkle, Vinca
Vitis Sp.*	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentos	Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants.

\*\*Do not use on container grown plants.

## STORAGE AND DISPOSAL

### PESTICIDE STORAGE

Do not contaminate water, food or feed by storage or disposal. Do not store in or around the home.

### PESTICIDE DISPOSAL

Wastes resulting from the use of this product must be disposed of on site or at an approved waste disposal facility.

### CONTAINER HANDLING

(For 50, 1000, 2000 pound bags and totes:)

Non-refillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then dispose of empty container in a sanitary land fill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

(For Killebrews and other bulk transport vehicles:)

Refillable container: Container handling instructions are not required. Cleaning before refilling is the responsibility of the refiller.

## CONDITIONS OF SALE AND WARRANTY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitation of Liability before using this product. If the terms are not acceptable, return the unopened container at once. By using this product, user and buyer accept the following Conditions, Disclaimer of Warranties, and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Harrell's, Inc. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** There are no warranties express or implied, of merchantability or of fitness for a particular purpose or otherwise, which extend beyond the statements made on this label. No agent of Harrell's, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, Harrell's, LLC disclaims any liability whatsoever for special, incidental, or consequential damages resulting from the use or handling of this product.

**LIMITATION OF WARRANTIES:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price, or at Harrell's, LLC option, the replacement of the product.

Manufactured and Guaranteed By:

Harrell's, LLC  
P. O. Box 807  
Lakeland, FL 33802

Growing a Better World® is a registered trademark of Harrell's, LLC

BARRICADE® is a registered trademark of Syngenta, Inc

EPA Accepted: 10/24/2017