EBANON

STARTER FERTILIZER

16-25-12 MESA

GUARANTEED ANALYSIS	GUARANTEED MINIMUM ANALYSIS
Total Nitrogen (N)	Total Nitrogen (N)
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	2.1% Other Water Soluble Nitrogen* Available Phosphate (P ₂ 0 ₅)
2.2% Combined Sulfur (S) Derived From: Ammonium Phosphate, Ammonium Sulfate,	2.2% Combined Sulfur (S) Chlorine (CI) (maximum)
Methylene Ureas, Urea, Muriate of Potash.	Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.
*3.3% Slowly available Nitrogen from Methylene Ureas.	*3.3% Slowly available Nitrogen from Methylene Ureas.
Chlorine (CI) not more than 8.0%	
U.S.A	CANAD

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

WARNING

Hazard Statements Causes skin irritation. Causes eye irritation. May be harmful if inhaled. May cause respiratory irritation.

Application Rates:

Spreader

Gandy

Spyker

Vicon

Lebanon Turf

Andersons AccuPro

Lesco (letter/numeric) K/22

Earthway Rotary

PennMulch HVO

ProScape SS

1,000 sq. ft. or 174 lbs. per acre.

To feed at the rate of 1.25 lb. Phosphate (P_2O_3) per 1,000 sq. ft., apply this product at 5.0 lbs per 1,000 sq. ft. or 260 lbs. per acre. To feed at the rate of 1lb. Phosphate

SUGGESTED SPREADER SETTINGS

1.25#P₂O₅/1,000 ft² 1#P₂O₅ /1,000 ft²

16

6 II

J/21

41/2

26

 (P_2O_5) per 1,000 sq. ft., apply this product at 4 lbs per

0

18

30

61/2 II

n 51/2

30

different settings for desired coverage.

contains phosphorus and may not be used for lawn

total nitrogen per 1,000 sq. ft. annually.

York or Virginia except when

before the application

of Maryland.

These settings were calibrated and field tested.

However, age and condition of spreader, speed of

operation and evenness of terrain may require slightly

Apply only to actively growing turf. Do not exceed 3.2 lbs. of

maintenance on turf in the states of Maryland, New Jersey, New

1) Providing nutrients to specific soils and target vegetation as

determined to be necessary in accordance with a soil test that

was conducted by a laboratory* performed no more than 3 years

accordance with the recommended application rates established

under Section 8-803.7 of the Agriculture Article, Annotated Code

2) Establishing vegetation for the first time, such as after land

*In Maryland, the soil testing laboratory must be identified

In New Hampshire: This fertilizer contains phosphate. In New

Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

disturbance, provided the application is conducted in

3) Re-establishing or repairing a turf area.

Precautionary Statements for handling Wash hands and exposed skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water. IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. If skin irritation occurs: Get medical advice/ attention. If eye irritation persists: Get medical advice/attention.

Store locked up. Store in a well-ventilated place. Keep container tightly closed.

Dispose of contents/container properly as required by law for waste

Keep out of reach of children Dispose in accordance with all federal, state and local regulations.

tions in your approved nutrient management plan.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention. In Maryland, New Jersey, New York and Virginia: This fertilizer Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommenda-

> Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.5 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

To feed at the rate of 0.60 kg Phosphate (P205) Equivalent to 2.5 kg of product per 100 sq. meters.

Application Rates:

To feed at the rate of 0.48 kg Phosphate (P_2O_5) Equivalent to 1.9 kg of product per 100 sq. meters.
SUGGESTED SPREADER SETTINGS

3066E3	IED SPREADER	i SETTINGS
Spreader 0.60 k	g P ₂ O ₂ /100 m ²	0.48 kgP ₂ 0 ₅ /100 n
Lebanon Turf	5 ½	41/2
Andersons AccuPro	0	M
Earthway Rotary	18	16
Gandy	30	28
Lely	6½ II	6 II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	0	N
ProScape SS	0	N
Spyker	5 ½	41/2
Vicon	30	26
Those cottings were	a a librated on	d field teeted

However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

800-233-0628 • (717) 273-1685



Manufactured by: Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042 800-233-0628 • (717) 273-1685 www.LebanonTurf.com

STARTER FERTILIZER

16-25-12 MESA

GUARANTEED ANALYSIS

Total Nitrogen (N) 1	16.0%
11.8% Ammoniacal Nitrogen	
1.2% Water Insoluble Nitrogen*	
0.9% Urea Nitrogen	
2.1% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ 0 ₅)	25.0%
Soluble Potash (K ₂ 0)	12.0%
Sulfur (S)	2.2%
2.2% Combined Sulfur (S)	

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

*3.3% Slowly available Nitrogen from Methylene Ureas.

GUARANTEED MINIMUM ANALYSIS

Total Nitrogen (N)	16.0%
11.8% Ammoniacal Nitrogen	
1.2% Water Insoluble Nitrogen*	
0.9% Urea Nitrogen	
2.1% Other Water Soluble Nitrogen*	
Available Phosphate (P ₂ 0 ₅)	25.0%
2.1% Other Water Soluble Nitrogen* Available Phosphate (P ₂ 0 ₅) Soluble Potash (K ₂ 0) Sulfur (S)	
Sulfur (S)	2.2%
0.00/ 0.11 1.01/ /0)	
Chlorine (CI) (maximum)	8.0%

Derived From: Ammonium Phosphate, Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

*3.3% Slowly available Nitrogen from Methylene Ureas.

CANADA

Lebanon brand fertilizers are formulated for professional use on turf, including golf courses, sports fields, and landscapes. Specific application rates and frequencies are determined by the professional applicator based on local soils, regulations, and climate.

Application Rates:

To feed at the rate of 1.25 lb. Phosphate $(P_2 0_5)$ per 1,000 sq. ft., apply this product at 5.0 lbs per 1,000 sq. ft. or 260 lbs. per acre. To feed at the rate of 1lb. Phosphate $(P_2 0_5)$ per 1,000 sq. ft., apply this product at 4 lbs per 1,000 sq. ft. or 174 lbs. per acre.

SUGGESTED SPREADER SETTINGS		
Spreader 1.25#	P,0,/1,000 ft ²	1#P,0,/1,000 ft2
Lebanon Turf	5½	4½
Andersons AccuPro	0	M
Earthway Rotary	18	16
Gandy	30	28
Lely	6½ II	6 II
Lesco (letter/numeric)	K/22	J/21
PennMulch HVO	0	N
ProScape SS	0	N
Spyker	51/2	41/2
Vicon	30	26
These settings were calibrated and field tested		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Apply only to actively growing turf. Do not exceed 3.2 lbs. of total nitrogen per 1,000 sq. ft. annually.

In Maryland, New Jersey, New York and Virginia: This fertilizer contains phosphorus and may not be used for lawn maintenance on turf in the states of Maryland, New Jersey, New York or Virginia except when

- 1) Providing nutrients to specific soils and target vegetation as determined to be necessary in accordance with a soil test that was conducted by a laboratory* performed no more than 3 years before the application
- Establishing vegetation for the first time, such as after land disturbance, provided the application is conducted in accordance with the recommended application rates established by the State
- 3) Re-establishing or repairing a turf area.
- *In Maryland, the soil testing laboratory must be identified under Section 8-803.7 of the Agriculture Article, Annotated Code of Maryland.

In New Hampshire: This fertilizer contains phosphate. In New Hampshire, phosphate fertilizers may only be used for turf maintenance in golf courses, parks, athletic fields, and sod farms.

WARNING

Hazard Statements

Causes skin irritation.
Causes eye irritation.
May be harmful if inhaled.

May cause respiratory irritation.

Precautionary Statements for handling

Wash hands and exposed skin thoroughly after handling.

Wear protective gloves/protective clothing/eye protection/face protection. If in eyes, rinse with water and seek medical attention if symptoms persist. Take off contaminated clothing and wash before reuse.

IF ON SKIN: Wash with plenty of soap and water.

IF INHALED: Call » POISON CENTER or doctor/physician if you feel unwell. If skin irritation occurs: Get medical advice/ attention.

If eye irritation persists: Get medical advice/attention.

Store locked up.

Store in a well-ventilated place.

Keep container tightly closed.

Dispose of contents/container properly as required by law for waste pesticides.

Keep out of reach of children

Precautionary Statements for disposal -

Dispose in accordance with all federal, state and local regulations.

KEEP OUT OF REACH OF CHILDREN.

Fertilizer may cause irritation of eyes, nose, throat, and skin. In case of contact with skin or eyes, flush with plenty of water; for eyes get medical attention.

Use in accordance with recommendations of a qualified individual or institution, such as, but not limited to a certified crop advisor, agronomist, university crop extension publication, or apply according to recommendations in your approved nutrient management plan.

Do not apply near water, storm drains or drainage ditches. Do not apply if heavy rain is expected. Apply this product only to your lawn, and sweep any product that lands on the driveway, sidewalk, or street back on to your lawn.

WARRANTY

NOTE To the extent consistent with applicable law, Buyer assumes all responsibility for safety and performance if this product is not used according to the directions.

Application Rates and Frequencies in Canada: Follow your professional landscape management plan for your local course or area. A suggested application rate is 1 to 2.5 Kg for 100 m², applied every 4 to 8 weeks during the local growing season.

Application Rates:

To feed at the rate of 0.60 kg Phosphate (P $_20_5$) Equivalent to 2.5 kg of product per 100 sq. meters.

To feed at the rate of 0.48 kg Phosphate (P205)
Equivalent to 1.9 kg of product per 100 sg. meters.

Equivalent to 110 kg of product por 100 oq. motoror			
SUGGESTED SPREADER SETTINGS			
Spreader 0.60 kg	g P ₂ O ₂ /100 m ²	0.48 kgP ₂ O ₅ /100 m ²	
Lebanon Turf	5 ½	41/2	
Andersons AccuPro	0	M	
Earthway Rotary	18	16	
Gandy	30	28	
Lely	6½ II	6 II	
Lesco (letter/numeric)	K/22	J/21	
PennMulch HVO	0	N	
ProScape SS	0	N	
Spyker	51/2	41/2	
Vicon	30	26	

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.



Information regarding the contents and levels of metals in this product is available on the Internet at: http://www.regulatory-info-lebsea.com

800-233-0628 • (717) 273-1685