



# Signature™ XTRA Stressgard®

A plant health promoting systemic fungicide that alleviates stress from both abiotic and biotic sources.

**ACTIVE INGREDIENT:**

Aluminum tris (O-ethyl phosphonate): .....60.00%

**OTHER INGREDIENTS:**.....40.00%

**TOTAL:** ..... 100.00%

EPA Reg. No. 432-1541

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

For MEDICAL and TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577  
For PRODUCT USE Information Call 1-800-331-2867

**Specimen Label**

432-1541 150420AV1

<b>FIRST AID</b>	
<b>If swallowed:</b>	<ul style="list-style-type: none"> <li>• Call a poison control center or doctor immediately for treatment advice.</li> <li>• Do not induce vomiting unless told to do so by a poison control center or doctor.</li> <li>• Have person sip a glass of water if able to swallow.</li> <li>• Do not give anything to an unconscious person.</li> </ul>
<b>If in eyes:</b>	<ul style="list-style-type: none"> <li>• Hold eyes open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing.</li> <li>• Call a poison control center or doctor for treatment advice.</li> </ul>
<p><b>Have the product container or label with you when calling a poison control center or doctor or going for treatment. In case of medical emergency for additional information call toll free 1-800-334-7577.</b></p>	
<p><b>NOTE TO PHYSICIAN:</b> Treat symptomatically.</p>	

## PRECAUTIONARY STATEMENTS

### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

#### CAUTION

- Harmful if swallowed.
- Causes moderate eye irritation.
- Avoid contact with eyes or clothing.
- Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

#### Personal Protective Equipment (PPE)

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions exist for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Applicators and other handlers must wear:

- Long-sleeved shirt
- Long pants
- Shoes plus socks
- Chemical-resistant (such as natural rubber) gloves (if more options are needed, follow the instructions for category A on an EPA chemical-resistance category selection chart).

#### Engineering Controls:

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

## USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

### ENVIRONMENTAL HAZARDS

Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters.

### CONDITIONS OF SALE AND LIMITATIONS OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions, Disclaimer of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Ineffectiveness, plant injury, other property damage, as well as other unintended consequences may result because of factors beyond the control of Bayer CropScience LP. Those factors include, but are not limited to, weather conditions, presence of other materials or the manner of use or application. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP MAKES NO OTHER WARRANTIES, EXPRESS OR IMPLIED, OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE OR OTHERWISE, THAT EXTEND BEYOND THE STATEMENTS MADE ON THIS LABEL. No agent of Bayer CropScience LP is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, BAYER CROPSCIENCE LP DISCLAIMS ANY LIABILITY WHATSOEVER FOR SPECIAL, INCIDENTAL OR CONSEQUENTIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT.

**LIMITATIONS OF LIABILITY:** TO THE EXTENT CONSISTENT WITH APPLICABLE LAW THE EXCLUSIVE REMEDY OF THE USER OR BUYER FOR ANY AND ALL LOSSES, INJURIES OR DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, WHETHER IN CONTRACT, WARRANTY, TORT, NEGLIGENCE, STRICT LIABILITY OR OTHERWISE, SHALL NOT EXCEED THE PURCHASE PRICE PAID, OR AT BAYER CROPSCIENCE LP'S ELECTION, THE REPLACEMENT OF PRODUCT.

### DIRECTIONS FOR USE

**It is a violation of federal law to use this product in a manner inconsistent with its labeling.**

**Read entire label before using this product.**

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

**PPE required for early entry to treated areas (that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water), is:**

- coveralls over long-sleeved shirt and long pants,
- socks and shoes,
- chemical-resistant gloves made of any of any waterproof material such as natural rubber  $\geq$  14 mils.

### NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard (WPS) for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses. Do not enter allow other to enter the treated area until sprays have dried.

### PRODUCT INFORMATION

**Signature™ Xtra Stressgard®** is a plant health promoting systemic fungicide that alleviates stress from both abiotic and biotic sources.

**For use on:**

- Turf on golf courses, sod farms, sport fields, institutional, municipal, and commercial sites.

### PLANT HEALTH

**Signature Xtra Stressgard** when applied at the prescribed rates and intervals promotes healthier roots, increases plant drought tolerance, and improves plant water utilization. This leads to a reduction in the need for hand watering and a general saving of water use. The resulting improvement in function of these healthier roots leads to more efficient uptake of plant nutrients improving cell turgidity and results in a healthier and sturdier plant structure reducing the impact of traffic or mechanical damage. These effects are consistent with the properties of anti-oxidants. A property of **Signature Xtra Stressgard** is to act as an inducer of systemic resistance in plants resulting in production of plant metabolites responsible for maintaining plant processes.

Cool season grasses such as tall fescues, bluegrasses and bentgrasses are prone to damage during summer heat and summer decline leading to reduced root mass in the summer months. Turf treated with **Signature Xtra Stressgard** retains turf quality through summer stress by alleviating plant stresses, enhancing disease resistance through induced system resistance and protecting the plant from the adverse effects of solar radiation.

Warm season grasses such as bermudagrass and zoysiagrass, influenced by the effects of insufficient sunlight, are prone to losing chlorophyll and premature senescence. **Signature Xtra Stressgard** reduces the negative effects of low light conditions, supports earlier spring green up, turf fill in and extends turf quality.

**Signature Xtra Stressgard** when applied in a program with turf growth regulators further enhances the overall quality of the turf.

### APPLICATION INFORMATION

**Signature Xtra Stressgard** must be applied as a foliar spray using ground application equipment. Apply in 1 to 5 gallons of spray solution per 1000 sq ft. This product may cause staining. Avoid off-target sites such as sidewalks, patios, driveways, pavers, or similar materials. Do not enter treated areas until spray has dried. Rinse spray equipment thoroughly with warm water and detergent.

### SPRAY MIXING AND COMPATIBILITY

**Signature Xtra Stressgard** may not be chemically compatible with foliar fertilizers or with copper-based fungicides and phytotoxicity may occur. To determine the physical compatibility of **Signature Xtra Stressgard** with any other product, use a small container to mix a small amount (e.g. 1 pint) of spray solution, containing all ingredients in the same order and ratio as the anticipated use. If any indication of physical incompatibility develops, do not use this mixture for spraying. Indication of incompatibility usually appears within 5 - 15 minutes after mixing. Read and follow all directions and precautions on this label and on the labels of any products for which a tank mixture is being considered.

### PRODUCT USE RESTRICTIONS

- Do not graze animals on treated turf.
- Do not feed clippings from treated turf to livestock and poultry.

### TURF USE DIRECTIONS

**Signature Xtra Stressgard** is a systemic fungicide, which may be used, in a seasonal program for the control of Pythium diseases, such as blight and root rot, and yellow tuft on common turfgrasses on golf courses, sod farms, sport fields, institutional, municipal, and commercial sites. Apply as a foliar spray, using 1 to 5 gallons of spray solution per 1,000 square feet, as indicated in the table below. Apply with a properly calibrated sprayer. Maintain agitation during spray operations.

### PLANT HEALTH

**Signature Xtra Stressgard** has a plant health component that is used for the prevention or alleviation of stress from solar radiation and other oxidative processes on turfgrasses. Begin preventative applications prior to the occurrence of adverse weather conditions and repeat at prescribed intervals. Use higher rates prior to the onset of severe stress. **Signature Xtra Stressgard** does not replace the use of soil amending adjuvants for soil hydrophobicity and must be used in conjunction with best turf management practices.

STRESS MANAGEMENT	RATE OZ PRODUCT / 1000 SQ FT	INTERVAL OF APPLICATIONS (DAYS)	APPLICATION DIRECTIONS
Stresses such as <ul style="list-style-type: none"> <li>• high solar radiation</li> <li>• high wilt potential</li> <li>• winter cold injury*</li> <li>• hot and humid weather</li> <li>• traffic wear</li> <li>• physical or mechanical damage</li> <li>• injury from the use of turf growth regulators*</li> <li>• injury from disease</li> <li>• reduced mowing heights</li> </ul>	2 - 4	7 - 14	Extend application interval beyond 14 days when climatic conditions are less favorable for turf growth. Do not mow or irrigate treated area until sprays have completely dried.
	4 - 6	14 - 21	
*Apply after damage has occurred			

#### APPLICATIONS FOR TURF DISEASES

DISEASE	RATE OZ PRODUCT / 1000 SQ FT	INTERVAL OF APPLICATIONS (DAYS)	APPLICATION DIRECTIONS
Pythium diseases, Yellow tuft	2 - 4	7 - 14	Begin preventive foliar spray application when conditions first favor disease and repeat at prescribed interval. Use the highest rate for the longer interval. Do not mow or irrigate treated area until sprays have completely dried.
	4 - 6	14 - 21	
	2 - 4	7	Use the more frequent interval when severe symptoms appear.
Bacterial Wilt	6	14	

#### USE AS A RESISTANCE MANAGEMENT TOOL

Fosetyl-AI has no history of resistance development in plant pathogenic fungi; include **Signature Xtra Stressgard** as a mixture or rotation partner in programs to manage resistance for fungicides with a high risk of resistance to labeled diseases.

#### TANK MIXTURES

A **Signature Xtra Stressgard** program encourages the development of a thick, vigorous turf stand that is better able to tolerate various stresses (heat, humidity, drying winds, traffic etc) that can have a negative effect on turf health. A thick, vigorous **Signature Xtra Stressgard** treated turf is better able to resist disease development and suppress the development of other pests such as weeds and algae.

**Summer Decline:** For control of Summer Decline caused by a complex of Pythium and Rhizoctonia diseases, **Signature Xtra Stressgard** can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. Do not exceed label dosage rates. This product cannot be mixed with any product containing a label prohibition against such mixing.

**Anthracnose:** For improved control of anthracnose **Signature Xtra Stressgard** can be tank-mixed with the products listed below in accordance with the most restrictive of label limitations and precautions. A preventative, multi-application program must be started prior to the appearance of the disease and the conditions favorable to anthracnose.

PRODUCT	DISEASE
Signature Xtra Stressgard + 26GT® or Interface® StressGard or Tartan® StressGard or Mirage® StressGard or Daconil Ultrex® or Daconil Action® or Fore® Rainshield™	Summer Decline (Pythium and Rhizoctonia spp.) and/or Anthracnose (Colletotrichum graminicola)
Signature Xtra Stressgard + Interface Stressgard or Fore Rainshield or Daconil Ultrex	Leaf spots Bermudagrass Decline

### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

#### PESTICIDE STORAGE

Store in original container and keep tightly closed when not in use. Store in a cool dry place. Avoid cross-contamination with other pesticides.

#### PESTICIDE DISPOSAL

Pesticides wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be used according to label instructions, contact your State Pesticide or Environmental Control Agency or Hazardous Waste representative at the nearest EPA regional office for guidance in proper disposal methods.

#### CONTAINER HANDLING

##### **Rigid Non-refillable Containers that are too large to shake (i.e., with capacities greater than 5 gallons or 50 lbs)**

Non-refillable container. Do not reuse or refill this container. Refer to Bottom Discharge IBC or Top Discharge IBC, Drums, Kegs information as follows.

##### **Bottom Discharge IBC (e.g. – Schuetz Caged IBC or Snyder Square Stackable)**

Pressure rinsing the container before final disposal is the responsibility of the person disposing of the container. To pressure rinse the container before final disposal, empty the remaining contents from the IBC into application equipment or mix tank. Raise the bottom of the IBC by 1.5 inches on the side which is opposite of the bottom discharge valve to promote more complete product removal. Completely remove the top lid of the IBC. Use water pressurized to at least 40 PSI to rinse all interior portions. Continuously pump or drain rinsate into application equipment or rinsate collection system while pressure rinsing. Continue pressure rinsing for 2 minutes or until rinsate becomes clear. Replace the lid and close bottom valve.

Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

##### **Top Discharge IBC, Drums, Kegs (e.g.– Snyder 120 Next Gen, Bonar B120, Drums, Kegs)**

Triple rinsing the container before final disposal is the responsibility of the person disposing of the container. To triple rinse the container before final disposal, empty the remaining contents from this container into application equipment or mix tank. Fill the container at least 10 percent full with water. Agitate vigorously or recirculate water with the pump for 2 minutes. Rinse all interior surfaces. Pour or pump rinsate into application equipment or rinsate collection system. Repeat this procedure two more times. Once container is rinsed, offer for recycling if available or puncture and dispose of in a sanitary landfill or by incineration.

**Non-Seed Treatment Products in Non-Refillable Fiber Drums with Liners** Non-refillable container. Do not reuse or refill this container. Completely empty liner by shaking and tapping sides and bottom to loosen clinging particles. Empty residue into application equipment, then offer for recycling if available or dispose of in a sanitary landfill or by incineration. If drum is contaminated and cannot be reused, dispose of it in the manner required for its liner.

**Non-Seed Treatment Products in Non-Refillable Foil outer pouches of Water soluble Packets (WSP)** Offer foil pouch for recycling if available or dispose of empty pouch in the trash as long as WSP is unbroken.

##### **Rigid Non-Refillable containers with capacities smaller or equal to 5 gallons**

#### PLASTIC CONTAINERS:

Non refillable container. Do not reuse or refill this container. Tripled rinse container (or equivalent) promptly after emptying.

#### **SOLID Dillutable formulations:**

Triple rinse as follows: Empty the remaining contents into application equipment or a mix tank. Fill the container ¼ full with water and recap. Shake for 10 seconds. Pour rinsate into application equipment or a mix tank or store rinsate for later use or disposal. Drain for 10 seconds after the flow begins to drip. Repeat this procedure two more times. Then offer for recycling if available or puncture and dispose of in a sanitary landfill, or by incineration.

##### **Non-Seed Treatment Products in Non-Rigid, Non-refillable Containers = BAGS**

Nonrefillable container. Do not reuse or refill this container. Completely empty container into application equipment. Then offer for recycling if available or dispose of in a sanitary landfill or by other procedures approved by state and local authorities.

Bayer (reg'd), the Bayer Cross (reg'd), 26GT®, Interface®, Mirage®, Stressgard®, Tartan® and Signature™ are trademarks of Bayer. Daconil® and Daconil Ultrex® are registered trademarks of GB Biosciences Corporation. Fore® and Rainshield® are trademarks of Dow Agrosciences LLC.

Backed by  
**BAYER**

Produced for:  
Bayer Environmental Science  
A Division of Bayer CropScience LP  
2 T. W. Alexander Drive  
Research Triangle Park, NC 27709

Bayer